









Supplementary Materials







Supplementary Materials



Supplementary Materials


Supplementary Figure 1. Trends of incidence rates among each male Asian American racial/ethnic group by cancer site from 1990-2008. Racial/ethnic groups are represented by the following plot symbols: Asian Indian/Pakistani: Solid red squares, Chinese: solid blue squares, Filipino: open green squares, Japanese: open orange squares, Kampuchean: solid purple circles, Korean: solid grey circles, Laotian: open black circles, Vietnamese: solid cyan triangles. Data above is shown by cancer site: **A)** prostate, 95% CIs are: Asian Indian/Pakistani segment 1 0.3 to 4.1, segment 2 -29.5 to 12.2, segment 3 -11.1 to 38.8; Chinese -0.7 to 1.2; Filipino segment 1 4.5 to 35.4, segment 2 -25.8 to 13.6, segment 3 -1.7 to 0.7; Japanese segment 1 -16.9 to 58.5, segment 2 -24.8 to 0.3, segment 3 -6.7 to 24.8, segment 4 -4.9 to 1.3; Kampuchean -3.8 to 14.2; Korean 1.8 to 4.0; Laotian -6.8 to 11.1; Vietnamese -1.2 to 2.4; **B)** colon and rectum, 95% CIs are: Asian Indian/Pakistani -1.5 to 1.7; Chinese -2.6 to -1.2; Filipino -0.5 to 0.9; Japanese -1.3 to -0.4; Kampuchean -4.1 to 11.4; Korean 0.9 to 3.5; Laotian -0.4 to 17.1; Vietnamese -0.1 to 3.0; **C)** lung and bronchus, 95% CIs are: Asian Indian/Pakistani 0.4 to 5.3; Chinese segment 1 -7.8 to -0.6, segment 2 -0.8 to 1.3; Filipino -0.7 to 0.6; Japanese -0.7 to 0.9; Kampuchean -9.5 to 5.3; Korean -2.3 to 0.3; Laotian -3.9 to 0.6; Vietnamese -2.1 to 0.9; **D)** stomach, 95% CIs are: Asian Indian/Pakistani -5.4 to 4.3; Chinese -2.3 to -0.3; Filipino -3.7 to -0.8; Japanese -4.0 to -2.4; Kampuchean -27.0 to 36.5; Korean -1.9 to 0.8; Laotian -13.9 to 3.5; Vietnamese -4.4 to -1.1; **E)** liver, 95% CIs are: Asian Indian/Pakistani 0.2 to 4.4; Chinese -1.2 to 0.4; Filipino 0.4 to 2.7; Japanese -0.1 to 2.7; Kampuchean -0.9 to 6.3; Korean segment 1 0.4 to 3.7, segment 2 -44.7 to 19.1; Laotian -1.9 to 4.2; Vietnamese 0.3 to 2.8; **F)** bladder, 95% CIs are: Asian Indian/Pakistani -0.9 to 3.0; Chinese -1.1 to 1.8; Filipino -0.8 to 1.5; Japanese 1.1 to 3.2; Korean 0.9 to 6.2; Vietnamese -1.2 to 4.9; **G)** non-Hodgkin lymphoma, 95% CIs are: Asian Indian/Pakistani -2.0 to 3.7; Chinese -0.5 to 1.7; Filipino -0.4 to 1.4; Japanese -0.4 to 2.2; Kampuchean -15.3 to 27.3; Korean -1.0 to 4.8; Laotian -14.1 to 16.9; Vietnamese -2.6 to 2.7; and **H)** pancreas, 95% CIs are: Asian Indian/Pakistani -1.7 to 10.8; Chinese -2.0 to 0.8; Filipino -1.6 to 1.3; Japanese 1.1 to 3.5; Korean 2.0 to 5.9; Laotian -20.1 to 15.1; Vietnamese -2.4 to 3.5. An **asterisk** indicates the 95% CI for the APC does not include 0. † Joinpoint and observed rates are based on 2 year groups (1990-1991, 1992-1993, 1994-1995, 1996-1997,..., 2008). ‡ Joinpoint and observed rates are based on 5 year groups (1990-1994, 1995-1999, 2000-2004, 2005-2008). The sample sizes were too small to estimate trends for bladder cancer among Laotians and Kampuchean and pancreatic cancer among Kampuchean.

Supplementary Materials




Supplementary Materials


E: Stomach




F: Liver



G: Cervix



H: Thyroid



Supplementary Materials

Supplementary Figure 2. Trends of incidence rates among each female Asian American racial/ethnic group by cancer site from 1990-2008. Racial/ethnic groups are represented by the following plot symbols: Asian Indian/Pakistani: Solid red squares, Chinese: solid blue squares, Filipino: open green squares, Japanese: open orange squares, Kampuchean: solid purple circles, Korean: solid grey circles, Laotian: open black circles, Vietnamese: solid cyan triangles

Data above is shown by cancer site: **A)** breast, Asian Indian/Pakistani 2.4 to 3.6; Chinese 0.6 to 1.9; Filipino 0.9 to 1.7; Japanese segment 1 0.2 to 5.4, segment 2 - 4.1 to 0.6; Kampuchean -5.9 to 15.4; Korean 4.0 to 5.4; Laotian 0.0 to 6.4; Vietnamese 0.1 to 2.2; **B)** colon and rectum, Asian Indian/Pakistani -1.6 to 4.8; Chinese -1.3 to 0.0; Filipino 0.0 to 2.0; Japanese segment 1 -0.4 to 1.9, segment 2 -7.1 to -1.3; Kampuchean -7.8 to 23.9; Korean 1.7 to 3.9; Laotian 4.0 to 7.7; Vietnamese -0.4 to 2.5; **C)** lung and bronchus, Asian Indian/Pakistani -2.0 to 4.5; Chinese -1.1 to 0.5; Filipino 1.4 to 2.8; Japanese segment 1: 2.5 to 9.8, segment 2: -8.2 to 2.1, segment 3: -7.6 to 29.2, segment 4: -8.2 to 1.1; Kampuchean -6.1 to 6.2; Korean 0.6 to 3.6; Laotian -10.1 to 9.5; Vietnamese -2.3 to 0.6; **D)** uterus, Asian Indian/Pakistani 0.3 to 5.8; Chinese segment 1 -1.7 to 1.7, segment 2 1.4 to 12.9; Filipino 2.4 to 3.7; Japanese 0.1 to 2.0; Kampuchean -12.8 to 15.9; Korean -0.6 to 5.0; Laotian -8.4 to 2.6; Vietnamese -0.2 to 4.1; **E)** stomach, Asian Indian/Pakistani -4.2 to 7.8; Chinese -2.8 to -0.2; Filipino -1.5 to 1.3; Japanese -4.8 to -2.6; Kampuchean -28.8 to 28.1; Korean -0.8 to 1.4; Laotian -9.2 to -1.0; Vietnamese -6.2 to -0.3; **F)** liver, Asian Indian/Pakistani -3.3 to 13.6; Chinese -0.3 to 2.2; Filipino 0.2 to 3.9; Japanese 0.2 to 2.5; Kampuchean -9.2 to 22.2; Korean -2.6 to 0.9; Laotian -1.3 to 6.0; Vietnamese segment 1 -9.2 to 59.5, segment 2 -16.5 to 12.7, segment 3 -0.2 to 9.2; **G)** cervix, Asian Indian/Pakistani -3.2 to 1.2; Chinese -6.3 to -3.1; Filipino -5.0 to -2.2; Japanese -3.1 to -0.1; Kampuchean -18.3 to 16.8; Korean -6.0 to -3.2; Laotian -10.4 to -4.1; Vietnamese -9.5 to -7.4; and **H)** thyroid, Asian Indian/Pakistani 4.0 to 9.7; Chinese 2.3 to 5.4; Filipino 1.7 to 3.3; Japanese 0.5 to 4.0; Kampuchean -8.2 to 19.6; Korean; Laotian -5.2 to 0.7; Vietnamese -1.7 to 1.8. An **asterisk** indicates the 95% CI for the APC does not include 0. † Joinpoint and observed rates are based on 2 year groups (1990-1991, 1992-1993, 1994-1995, 1996-1997,..., 2008). ‡ Joinpoint and observed rates are based on 5 year groups (1990-1994, 1995-1999, 2000-2004, 2005-2008).